

## INVARIANT SPIN STRUCTURES ON RIEMANN SURFACES

SADOK KALLEL AND DENIS SJERVE \*

ABSTRACT. We investigate the action of the automorphism group of a closed Riemann surface on its set of theta characteristics (or spin structures). We give criteria for when an automorphism fixes all spin structures, or when it fixes just one. The case of hyperelliptic curves and of the Klein quartic are discussed in detail.

## 1. INTRODUCTION

Spin structures on Riemann surfaces or “theta characteristics” are classical objects of great use and interest in all of geometry, topology and physics. Let  $C$  be a closed Riemann surface and write  $U(C)$  for its unit tangent bundle. A spin structure on  $C$  is a cohomology class in  $H^1(U(C); \mathbb{Z}_2)$  whose restriction to each fiber is a generator of  $H^1(S^1; \mathbb{Z}_2)$ . There is of course a great number of ways to see what a spin structure is, some of which we will encounter in the course of this paper.

Every Riemann surface  $C$  has  $2^{2g}$  distinct spin structures where  $g$  is the genus of  $C$ . Generally the set of spin structures  $Spin(C)$  is an affine space associated with  $H^1(C; \mathbb{Z}_2)$  and hence there is a (non-canonical)  $1 - 1$  correspondence between  $Spin(C)$  and  $H^1(C; \mathbb{Z}_2)$ .

The group of automorphisms  $Aut(C)$  acts on  $H^1(U(C); \mathbb{Z}_2)$  and hence on  $Spin(C)$  by pullback. This action and the action on  $H^1(C; \mathbb{Z}_2)$  are however not compatible under the above  $1 - 1$  correspondence, but are closely related. They differ by a term involving the intersection pairing.

In this paper we study invariant spin structures on compact Riemann surfaces. Motivation for this work comes from a question by Jack Morava to the authors and from a paper of Atiyah [3]. In that paper Atiyah proves that if  $f : C \rightarrow C$  is an automorphism of a compact Riemann surface  $C$  then there necessarily exists a spin structure  $\xi$  that is invariant under  $f : C \rightarrow C$ .

This raises natural questions.

- (1) How many spin structures are invariant under  $f : C \rightarrow C$ ?
- (2) How large can the isotropy subgroup of a given spin structure in  $Aut(C)$  be?
- (3) Is there an automorphism  $f : C \rightarrow C$  that leaves only one spin structure invariant?
- (4) Is there a non-trivial automorphism  $f : C \rightarrow C$  that leaves every spin structure invariant?

We will give complete answers to these questions and more. The following is an answer to (4) and shows how restrictive that question is.

**Theorem 1.1.** *A non-trivial automorphism  $f : C \rightarrow C$  leaves every spin structure invariant if, and only if,  $C$  is hyperelliptic and  $f$  is the hyperelliptic involution.*

In fact one can determine precisely the number of invariant spin structures for an automorphism  $f$  in terms of its symplectic action on homology. Let  $A \in SL_{2g}(\mathbb{Z})$  denote the  $2g \times 2g$  matrix representing the induced isomorphism  $f_* : H_1(C; \mathbb{Z}) \rightarrow H_1(C; \mathbb{Z})$ , with respect to some basis, and let  $\bar{A}$  is its mod 2 reduction. If  $h$  is the dimension of the eigenspace of  $\bar{A}$  associated to the eigenvalue 1, then the number of  $f$ -invariant spin structures is  $2^h$ . This is our answer to (1).

---

\*Research partially supported by NSERC Discovery grant A7218.

Note that if  $f$  has order  $n$ , then necessarily  $h \geq 2k$  where  $k$  is the genus of the orbit surface  $C/\mathbb{Z}_n$ . This is because  $2k$  is the dimension of the eigenspace of  $A$  associated to the eigenvalue 1.

In contrast with theorem 1.1, by analyzing the similarity class of  $A$  in  $SL_{2g}(\mathbb{Z})$  we obtain

**Theorem 1.2.** *Suppose  $f : C \rightarrow C$  is an automorphism of order  $n$ , where  $n$  is odd. Then  $f$  leaves only one spin structure invariant if, and only if, the associated orbit surface  $C/\mathbb{Z}_n$  has genus zero.*

The above theorems are based on a short calculation that we give in section 2 using constructions of Johnson [8]. Let  $A, \bar{A}$  be as above, and let  $X = (x_1, \dots, x_{2g})^T$  be a column vector representing an element of  $H^1(C; \mathbb{Z})$ , and let  $\bar{X}$  be its mod 2 reduction. Then

**Proposition 1.3.** *The spin structures left invariant by an automorphism  $f : C \rightarrow C$  are in 1-1 correspondence with solutions  $\bar{X}$  of the matrix equation  $(\bar{A}^T - I)\bar{X}^T = 0$ .*

In particular we see that an automorphism  $f$  of  $C$ , or more generally an orientation preserving diffeomorphism, fixes all spin structures if and only if  $f$  acts trivially on  $H_1(C; \mathbb{Z}_2)$ . This is one form of the main theorem A of Sipe [15].

Another immediate corollary of proposition 1.3 and of computations of [14] is that the Klein quartic curve  $\mathcal{K}$  of genus 3 and maximal group of automorphisms has a unique invariant spin structure. Describing the geometry behind this structure requires the description of spin structures in terms of divisor classes (cf. §5, Appendix).

Consider the points  $a = [1 : 0 : 0]$ ,  $b = [0 : 1 : 0]$  and  $c = [0 : 0 : 1]$  belonging to  $\mathcal{K}$ , where  $\mathcal{K}$  is described as the locus  $x^3y + y^3z + z^3x = 0$  in  $\mathbb{P}^2$ . Let  $K$  be the canonical divisor on  $\mathcal{K}$  which is determined by the intersection of the curve with any hyperplane  $\mathbb{P}^1 \subset \mathbb{P}^2$ . In §4 we prove:

**Theorem 1.4.** *The divisor class  $\theta := 2a + 2b + 2c - K$  is the unique spin structure on the Klein curve  $\mathcal{K}$  fixed by the entire group  $\text{Aut}(\mathcal{K}) = \text{PSL}(2, \mathbf{F}_7)$ .*

The existence of a unique spin structure on Klein's curve is well-known to algebraic geometers [6]. It comes as follows. Let  $X(p)$ ,  $p > 5$  prime, denote the modular curve defined as a compactification of the upper half-plane by the action of the principal congruence subgroup of level  $p$ . The group  $G = \text{PSL}(2, \mathbf{F}_p)$  acts as a group of automorphisms on  $X(p)$ . Then [1, 5] the group  $\text{Pic}(X(p))^G$  of invariant line bundles is infinite with cyclic generator a  $(2p - 12)$ -th root of the canonical bundle of degree  $(p^2 - 1)/24$ . When  $p = 7$ ,  $X(7) = \mathcal{K}$  and the generator is a square root of the canonical bundle hence the spin structure in question.

Note that if  $\text{Aut}(C, L)$  is the subgroup of all automorphisms of  $\text{Aut}(C)$  that fix a spin structure  $L$ , then the above corollary shows it is possible that  $\text{Aut}(C, L)$  reaches the maximum Hurwitz bound of  $84(g - 1)$ . This answers our question (2).

Even though the above basic results do not seem to exist in the literature, we make a cautious claim to originality since some of what we prove might be known to algebraic geometers, coated in some orthogonal language to ours. Complementary results of a similar nature as ours can be found in [15]. A relevant discussion from the point of view of mapping class groups is in [10]. Finally a growing number of references on this question have to do with moduli spaces of spin curves.

This note is organized as follows. In §2 we collect the background material we need and prove our main results. In §3 we give a complete count of invariant spin structures for automorphisms of hyperelliptic surfaces based on a combinatorial definition of spin structures due to Mumford. See propositions 3.3 and 3.4. In the last section §4 we discuss the case of quartics in general and the Klein curve in particular. An appendix clarifies the relationship between the two different definitions of spin structures we use in this paper.

## 2. SPIN STRUCTURES AND AUTOMORPHISMS

We assume  $C$  is smooth, closed, connected and orientable. Let  $S^1 \xrightarrow{i} U(C) \xrightarrow{\pi} C$  be the unit tangent bundle. We adopt Johnson's definition of a spin structure [8]; namely this is a cohomology class  $\xi \in H^1(U(C); \mathbb{Z}_2)$  that restricts to a generator of  $H^1(S^1; \mathbb{Z}_2) \approx \mathbb{Z}_2$  for every fiber  $S^1$ .

There are short exact sequences

$$\begin{array}{ccccccc} 0 & \rightarrow & H_1(S^1; \mathbb{Z}_2) & \xrightarrow{i_*} & H_1(U(C); \mathbb{Z}_2) & \xrightarrow{\pi_*} & H_1(C; \mathbb{Z}_2) \rightarrow 0 \\ 0 & \leftarrow & H^1(S^1; \mathbb{Z}_2) & \xleftarrow{i^*} & H^1(U(C); \mathbb{Z}_2) & \xleftarrow{\pi^*} & H^1(C; \mathbb{Z}_2) \leftarrow 0 \end{array}$$

from which one deduces that the set  $Spin(C)$  of spin structures on  $C$  is in  $1-1$  correspondence with  $H^1(C; \mathbb{Z}_2) \approx \mathbb{Z}_2^{2g}$ , and therefore there are  $2^{2g}$  spin structures on  $C$ . This correspondence is not a group isomorphism and  $Spin(C)$  is in fact the non-trivial coset of  $H^1(C; \mathbb{Z}_2)$  in  $H^1(U(C); \mathbb{Z}_2)$ .

Suppose  $\omega$  is a smooth simple closed curve on  $C$ . Let  $\bar{\omega}$  denote the lift of  $\omega$  to  $U(C)$  given by the unit tangent vector at each point of  $\omega$ . There are actually 2 lifts, but they are homotopic in  $U(C)$  by the homotopy that rotates one tangential direction to the other.

We will let  $z \in H_1(U(C); \mathbb{Z}_2)$  denote the class represented by the tangential framing on  $\partial(D^2)$ , where  $D^2$  is any closed 2-disc in  $C$ . As is nicely explained by Johnson, "Intuitively a spin structure  $\zeta \in Spin(C)$  is a function assigning a number mod 2 to each framed curve of  $C$ , subject to the usual homological conditions and also that the boundary of a disc in  $C$  tangentially framed receives one".

**Lemma 2.1.** (Johnson [8]) *Suppose the homology class  $u \in H_1(C; \mathbb{Z}_2)$  is represented by  $\omega_1 + \omega_2 + \dots + \omega_r$ , where  $\{\omega_1, \omega_2, \dots, \omega_r\}$  is a set of non-intersecting smooth simple closed curves. Then  $\bar{\omega}_1 + \bar{\omega}_2 + \dots + \bar{\omega}_r + rz$  depends only on the class  $u$ , and not on the particular representation.*

**Definition 2.2.**  $\tilde{u} = \bar{\omega}_1 + \bar{\omega}_2 + \dots + \bar{\omega}_r + rz$ .

This canonical lifting from  $H_1(C; \mathbb{Z}_2)$  to  $H_1(U(C); \mathbb{Z}_2)$  fails to be a homomorphism. This is made explicit by the following result in [8].

**Lemma 2.3.** *If  $a, b \in H_1(C; \mathbb{Z}_2)$  then  $\widetilde{a+b} = \tilde{a} + \tilde{b} + \langle a, b \rangle z$ , where  $\langle a, b \rangle$  is the intersection pairing.*

Let  $e_1, \dots, e_{2g}$  denote a basis of  $H_1(C; \mathbb{Z}) \approx \mathbb{Z}^{2g}$ . We do not assume that this basis is symplectic. The following is clear.

**Lemma 2.4.** *A basis for  $H_1(U(C); \mathbb{Z})$  is given by  $\tilde{e}_1, \tilde{e}_2, \dots, \tilde{e}_{2g}, z$ .*

Let the dual basis of  $H^1(U(C); \mathbb{Z})$  be denoted  $\zeta_1, \dots, \zeta_{2g}, \eta$ . Then the mod 2 reduction  $\bar{\eta} \in H^1(U(C); \mathbb{Z}_2)$  is a particular spin structure on  $C$ . It follows that the set of spin structures is given by

$$Spin(C) = \left\{ \sum_{i=1}^{2g} x_i \bar{\zeta}_i + \bar{\eta} \mid \text{all } x_i \in \mathbb{Z}_2 \right\}.$$

We can now determine the action of an automorphism  $f : C \rightarrow C$  on  $Spin(C)$ . Write  $f_*(e_i) = \sum_{j=1}^{2g} a_{ji} e_j$ ,  $i = 1, \dots, 2g$ , where the  $a_{i,j}$  are the entries of  $A \in SL_{2g}(\mathbb{Z})$ .

**Definition 2.5.**  $v_i = \sum_{1 \leq j_1 < j_2 \leq 2g} a_{j_1 i} a_{j_2 i} \langle e_{j_1}, e_{j_2} \rangle$ ,  $V = [v_1, v_2, \dots, v_{2g}]$ .

We use the notation  $V_f$  or  $V_A$  if we want to emphasize that the vector  $V$  comes from  $f$  or  $A$ . Note that if  $f$  is a diffeomorphism of  $C$  inducing  $f_*$  (resp.  $f^*$ ) on  $H_1$  (resp.  $H^1$ ) then  $\widetilde{f_*(a)} = f_*(\tilde{a})$  (and same for  $f^*$ ). This is because  $f_*(\bar{a}) = \overline{f_*(a)}$  since  $f_*$  acts by its differential

on the tangent space and because  $f_*(z) = z$ . The following easy computation is an immediate corollary of this fact

**Lemma 2.6.** *With  $f_*(e_i) = \sum_{j=1}^{2g} a_{ji}e_j$  for  $i = 1, \dots, 2g$ , we have*

$$\begin{aligned} f_*(\tilde{e}_i) &= \sum_{j=1}^{2g} a_{ji}\tilde{e}_j + v_i z \text{ and } f_*(z) = z \\ f^*(\zeta_i) &= \sum_{k=1}^{2g} a_{ik}\zeta_k \text{ and } f^*(\eta) = \sum_{k=1}^{2g} v_k\zeta_k + \eta. \end{aligned}$$

Thus the matrices for  $H_1(U(C); \mathbb{Z}) \xrightarrow{f_*} H_1(U(C); \mathbb{Z})$  and  $H^1(U(C); \mathbb{Z}) \xrightarrow{f^*} H^1(U(C); \mathbb{Z})$  are  $\begin{bmatrix} A & 0 \\ V & 1 \end{bmatrix}$  and  $\begin{bmatrix} A^T & V^T \\ 0 & 1 \end{bmatrix}$  respectively. A routine calculation gives

**Lemma 2.7.** *Suppose  $\xi = \sum_{i=1}^{2g} x_i \bar{\zeta}_i + \bar{\eta}$  is a spin structure. Then*

$$f^*(\xi) = \sum_{k=1}^{2g} \left( \sum_{i=1}^{2g} a_{ik}x_i + v_k \right) \bar{\zeta}_k + \bar{\eta}.$$

Let  $X$  denote the column vector  $(x_1, \dots, x_{2g})^T$ . Then

**Corollary 2.8.**  $\xi = \sum_{i=1}^{2g} x_i \bar{\zeta}_i + \bar{\eta}$  is an invariant spin structure for  $f : C \rightarrow C$  if, and only if,

$$(1) \quad (\bar{A}^T - I)\bar{X} = -\bar{V}^T$$

This is equivalent to  $(A^T - I)X \equiv -V^T \pmod{2}$ . Proposition 1.3 of §1 is now a consequence for if we suppose that  $\xi_1, \xi_2$  are invariant spin structures associated to column vectors  $X_1, X_2$ , then each vector satisfies equation (2.8), and therefore  $X = X_1 - X_2$  satisfies  $(\bar{A}^T - I)\bar{X}^T = 0$ .

**2.1. Proof of Main Theorems 1.1 and 1.2.** We first need information on the similarity class of the matrix  $A \in SL_{2g}(\mathbb{Z})$ .

**Definition 2.9.** Let  $\Phi_d(x)$  be the cyclotomic polynomial generated by the primitive  $d^{\text{th}}$  roots of unity and let  $C_d$  denote the  $\phi(d) \times \phi(d)$  companion matrix of  $\Phi_d(x)$ , that is

$$C_d = \begin{bmatrix} 0 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & 0 & \cdots & 0 \\ . & . & . & . & \cdots & 1 \\ -a_0 & -a_1 & . & . & \cdots & -a_{\phi(d)-1} \end{bmatrix}$$

where  $\phi(d)$  denotes Euler's totient function and  $\Phi_d(x) = a_0 + a_1x + \cdots + a_{\phi(d)-1}x^{\phi(d)-1} + x^{\phi(d)}$ .

Suppose  $f : C \rightarrow C$  is an automorphism of order  $n$ . If  $g \geq 2$  then  $A$  also has order  $n$ . It follows that over the rationals  $\mathbb{Q}$ ,  $A$  is similar to a direct sum of companion matrices. In fact, according to [16],  $\exists$  unique distinct divisors  $1 \leq d_1 < d_2 < \cdots < d_r \leq n$  and unique positive integers  $e_1, \dots, e_r$  such that

- $A$  is similar to  $e_1 C_{d_1} \oplus e_2 C_{d_2} \oplus \cdots \oplus e_r C_{d_r}$  as a matrix in  $GL_{2g}(\mathbb{Q})$ .
- $n = \text{LCM}(d_1, \dots, d_r)$  and  $2g = e_1 \phi(d_1) + e_2 \phi(d_2) + \cdots + e_r \phi(d_r)$ .

The minimal and characteristic polynomials of  $C_d$  are both  $\Phi_d(x)$ . It follows that the minimal and characteristic polynomials of  $A$  are

$$\mu_A(x) = \prod_{i=1}^r \Phi_{d_i}(x), \quad \gamma_A(x) = \prod_{i=1}^r \Phi_{d_i}(x)^{e_i} \text{ respectively.}$$

From the factorization  $x^{n-1} + x^{n-2} + \dots + 1 = \prod_{d|n, d>1} \Phi_d(x)$  it follows that  $n = \prod_{d|n, d>1} \Phi_d(1)$  and  $\det(I - A^T) = \gamma_A(1) = \prod_{i=1}^r \Phi_{d_i}^{e_i}(1)$ . The following lemma is well known.

**Lemma 2.10.**  $\Phi_d(1) = \begin{cases} 0 & \text{if } d = 1 \\ p & \text{if } d = p^k, p = \text{a prime} \\ 1 & \text{in all other cases} \end{cases}$

**Corollary 2.11.**  $\bar{A}^T - I$  is invertible (as a matrix over  $\mathbb{Z}_2$ ) if, and only if,  $d_1 > 1$  and none of the  $d_i$  are powers of 2.

From proposition (1.3) it follows that  $f : C \rightarrow C$  has a unique invariant spin structure if, and only if,  $\bar{A}^T - I \in SL_{2g}(\mathbb{Z}_2)$ . This happens if, and only if,  $\det(A - I)$  is odd. Now theorem (1.2) of §1 will follow from this observation.

**Theorem 2.12.** Suppose  $f : C \rightarrow C$  is an automorphism of order  $n$ , where  $n$  is odd. Then  $f$  leaves only one spin structure invariant if, and only if, the associated orbit surface  $C/\mathbb{Z}_n$  has genus zero.


*Proof.* If  $V$  is a vector space and  $L : V \rightarrow V$  is a linear map let's write  $E(L, \lambda) = \{v \in V \mid L(v) = \lambda v\}$  the eigenspace of  $L$  associated to  $\lambda$ . Let  $\mathcal{H}$  (resp.  $\mathcal{H}^*$ ) denote the complex vector space of holomorphic (resp. anti-holomorphic) differentials on the curve  $C$ ,  $\dim \mathcal{H} = g$  the genus of  $C$ . The automorphism  $f : C \rightarrow C$  induces linear automorphisms  $L : \mathcal{H} \rightarrow \mathcal{H}$ ,  $L^* : \mathcal{H}^* \rightarrow \mathcal{H}^*$ . The subspaces  $E(L, 1)$  and  $E(L^*, 1)$  correspond to the  $\mathbb{Z}_n$ -invariant differentials, where  $\mathbb{Z}_n$  is the cyclic subgroup generated by  $f$ , and thus they have dimension  $h := \text{genus}(C/\mathbb{Z}_n)$ . Now there is an isomorphism  $H^1(C; \mathbb{Z}) \otimes \mathbb{C} \approx \mathcal{H} \oplus \mathcal{H}^*$  which is compatible with  $f^* : H^1(C; \mathbb{Z}) \otimes \mathbb{C} \rightarrow H^1(C; \mathbb{Z}) \otimes \mathbb{C}$  on the one hand, and with  $L \oplus L^* : \mathcal{H} \oplus \mathcal{H}^* \rightarrow \mathcal{H} \oplus \mathcal{H}^*$  on the other. From this it follows that the dimension of  $E(f^*, 1) = E(A, 1)$  equals  $2h$ . From the rational canonical form  $e_1 C_{d_1} \oplus e_2 C_{d_2} \oplus \dots \oplus e_r C_{d_r}$  we see that the dimension of  $E(A, 1)$  is 0 if  $d_1 > 1$  and  $e_1$  otherwise. The result now follows from corollary 2.11 and proposition 1.3.  $\square$

Next we give a proof of theorem 1.1.

*Proof.* Suppose  $C$  is hyperelliptic and  $f$  is the hyperelliptic involution  $J$ . Then the induced isomorphism  $J_* : H_1(C; \mathbb{Z}) \rightarrow H_1(C; \mathbb{Z})$  is  $-I$ . Therefore  $V_J = [0, 0, \dots, 0]$  and equation (1) for an invariant spin structure becomes trivial.

Conversely, suppose  $f : C \rightarrow C$  is a non-trivial automorphism leaving every spin structure invariant. Then equation (1) is valid for all vectors  $X$  and therefore  $\bar{A} = I$ . By a theorem of Serre (see p. 275 of [7]) we see that  $A$  must have order 2, and therefore so does the automorphism  $f : C \rightarrow C$ . Now a theorem of Nielsen [13] states that  $f$  is determined up to conjugacy by its fixed point data. In the case of an automorphism of order 2 the fixed point data is just the number of fixed points, and therefore two involutions with the same number of fixed points are conjugate. Let the number of fixed points be  $r$ . It is known [7] that if  $f \neq id$ ,  $r \leq 2g + 2$ .

Assume  $r < 2g + 2$ . Let  $2s = 2g + 2 - r > 0$  ( $r$  must be even because of the Riemann-Hurwitz formula). Then  $f$  must be conjugate to an automorphism as depicted in the Figure 1, and

FIGURE 1. An involution with  $r < 2g + 2$  fixed points

therefore we can choose a basis so that

$$A = \begin{bmatrix} -I_r & 0 & 0 \\ 0 & 0 & -I_s \\ 0 & -I_s & 0 \end{bmatrix}$$

But this contradicts the equation  $\bar{A} = I$ , and therefore  $r = 2g + 2$ . Thus  $C$  is hyperelliptic and  $f$  is the hyperelliptic involution.  $\square$

**2.2. Corollaries.** Corollary 2.8 and the proof of theorem 1.1 have some elementary consequences. Let  $\rho : \text{Aut}(C) \longrightarrow Sp_{2g}(\mathbb{Z}_2)$  be the homomorphism induced from the action on  $H_1(C; \mathbb{Z}_2)$  where  $Sp_{2g}(\mathbb{Z})$  is the symplectic modular group and  $Sp_{2g}(\mathbb{Z}_2)$  its mod-2 reduction.

**Corollary 2.13.** [7]  *$\rho$  is a monomorphism if and only if  $C$  is non-hyperelliptic. In the hyperelliptic case  $\ker(\rho) = \mathbb{Z}_2$  is generated by the hyperelliptic involution.*

Let  $\Gamma_g(C)$  be the mapping class group of  $C$ ; that is the group of isotopy classes of orientation preserving diffeomorphisms of  $C$ . It is well-known that  $\Gamma_g(C)$  is a finitely generated group with generators the Dehn twists around representative loops of a symplectic basis of  $C$ . If  $[f] \in \Gamma_g$ , then the action of  $f$  on  $\text{Spin}(C)$  (viewed as cohomological classes) is independent of the choice of this representative so that  $\Gamma_g$  acts on  $\text{Spin}(C)$ . Let  $S_g \subset \Gamma_g$  be the subgroup of  $\Gamma_g$  that fixes all spin structures. The following is immediate

**Corollary 2.14.** [15] *The subgroup  $S_g$  is precisely the subgroup of elements that induce the identity on  $H_1(C; \mathbb{Z}_2)$ .*

In fact in [15], Sipe states her theorem for all  $n$ -roots and she identifies  $S_g$  with the subgroup of elements which induce the identity on  $H_*(U(C); \mathbb{Z}_2)$ . But an orientation preserving automorphism is the identity on  $H_*(U(C); \mathbb{Z}_2)$  if and only if it is the identity on  $H_*(C; \mathbb{Z}_2)$ .

### 3. SPIN STRUCTURES ON HYPERELLIPTIC CURVES

We use a convenient description of the spin structures on a hyperelliptic curve in terms of divisors due to Mumford ([12] or [2], appendix B) to count invariant spin structures for subgroups of the automorphism group.

From the appendix we have that

$$(2) \quad \text{Spin}(C) = \{D \in \text{Cl}(C) \mid 2D = K\}$$

where  $\text{Cl}(C)$  is the divisor class group of  $C$ . We will denote by  $J_2(C)$  the subgroup of points of order two in  $J(C) = \text{Cl}^0(C)$ , the “Jacobian” of line bundles of degree zero. See [2].

**3.1. Spin Divisors on a Hyperelliptic Curve.** We consider the hyperelliptic surface  $y^2 = \prod_{i=1}^{2g+2} (x - e_i)$  having genus  $g$  and branch set  $B = \{e_1, \dots, e_{2g+2}\}$ . The  $e_i$  are distinct points in the Riemann sphere. We will write  $\pi : C \longrightarrow \mathbb{P}^1$  for the degree two covering map sending  $(x, y) \mapsto x$ , and we let  $p_i$  be the ramification points such that  $\pi(p_i) = e_i$ . A description of the canonical divisor  $K$  of  $C$  is completely standard. Let  $D$  be any divisor of the form  $2p_i$  or  $x + y$  if  $x, y \notin \{p_1, \dots, p_{2g+2}\}$  (i.e.  $D$  is the divisor associated to the line bundle  $L$  obtained as the pullback of the unique line bundle over  $\mathbb{P}^1$  of degree  $+1$ ). Then [2, 12]

$$(3) \quad K_C = -2D + p_1 + \dots + p_{2g+2} = (g-1)D$$

We can next determine the divisor of an element  $L \in J_2(C)$ . Let  $\phi : \mathbb{P}^1 \longrightarrow \mathbb{C}$  be the meromorphic function  $\phi(z) = (z - e_i)/(z - e_j)$ ,  $i \neq j$ , and consider the composite  $C \xrightarrow{\pi} \mathbb{P}^1 \xrightarrow{\phi} \mathbb{C}$ . The divisor of the composite is  $(\phi \circ \pi) = 2p_i - 2p_j$  and hence by definition  $p_i - p_j$  is the divisor of a line bundle of order two on  $C$  ( $i \neq j$ ). Using the same argument with the rational function  $\phi(z) = (z - e_i)(z - e_j)/z^2$  shows that  $E = p_i - p_j = D - p_i - p_j$ . Generally one sees that if  $L \in J_2(C)$ , then it is represented by either one of the divisors

$$(4) \quad E = p_{i_1} + \dots + p_{i_k} - p_{j_1} - \dots - p_{j_k} = kD - p_{l_1} - \dots - p_{l_{2k}}$$

Note that the above formula shows that any finite set  $T \subset \{1, \dots, 2g+2\}$  of even cardinality gives an element in  $J_2(C)$ . One can further see that any such  $T$  and its complement  $T^c \subset B$  give rise to the same element. If we define  $E_g$  to be the quotient

$$E_g := \{T \mid T \subseteq B, |T| \text{ even}\} / \sim, \text{ where } T' \sim T \iff T' = T \text{ or } T' = T^c$$

Then the map

$$\alpha : E_g \longrightarrow J_2(C) \quad , \quad T \mapsto \alpha_T = \sum_{i \in T} (p_i - p_{2g+2})$$

defines an automorphism between two copies of  $\mathbb{Z}_2^{2g}$  [5].

**Theorem 3.1.** [12] *Let  $C$  be as above,  $p_i \in C$ . Then every theta characteristic is of the form*

$$\theta_T = kD + p_{i_1} + \dots + p_{i_{g-1-2k}}$$

for  $-1 \leq k \leq \frac{g-1}{2}$ ,  $T := \{p_{i_1}, \dots, p_{i_{g-1-2k}}\}$  with the  $i_\alpha$  distinct. Moreover such a representation is unique if  $k \geq 0$  and subject to a single relation  $-D + p_{i_1} + \dots + p_{i_{g+1}} = -D + p_{j_1} + \dots + p_{j_{g+1}}$  when  $k = -1$ .

*Proof.* An easy sketch is in [2], exercises 26-32, p. 287-288. We work in the divisor class group where we have the (equivalent) relations  $p_{i_1} + \dots + p_{i_{g+1}} = p_{j_1} + \dots + p_{j_{g+1}}$  and  $K = (g-1)D$ . With  $E$  as in the theorem we can then compute

$$(5) \quad 2E = 2kD + 2p_{i_1} + \dots + 2p_{i_{g-1-2k}} = 2kD + (g-1-2k)D = (g-1)D = K$$

which implies that  $E$  is a spin divisor. The previous paragraphs also show that

$$p_{i_1} + \dots + p_{i_r} = p_{j_1} + \dots + p_{j_r} \iff r = g+1$$

for distinct  $i_\alpha$ 's and  $j_\beta$ 's so that a representation of  $E$  as in (5) is unique when  $0 \leq k \leq (g-1)/2$ , thus yielding  $\sum_{k=0}^{(g-1)/2} \binom{2g+2}{g-1-2k}$  distinct spin structures. The one relation for  $r = g+1$  yields  $\frac{1}{2} \binom{2g+2}{g+1}$  additional structures. Since this adds up to exactly  $2^{2g}$ , we have accounted for all spin structures this way.  $\square$

Since to each branched point corresponds a unique ramification point, we can write

**Lemma 3.2.** *Spin( $C$ ) is the set of all  $T \subset B$  such that  $|T| \equiv g+1 \pmod{2}$  modulo the equivalence relation  $T \sim T^c$ . This has a natural affine structure over  $E_g$  given by  $\theta_S + \alpha_T = \theta_{T+S}$ .*

**3.2. Automorphisms.** Let  $f : C \longrightarrow C$  be an automorphism of a Riemann surface. Then  $f$  acts on  $\text{Spin}(C)$  as in (2) by sending

$$f(\sum n_i P_i) = \sum n_i f^{-1}(P_i)$$

Indeed one way to see this is to replace the cotangent bundle by the tangent bundle, and since  $f$  is an automorphism, replace the pullback  $f^* : T_{f(x)}^* C \longrightarrow T_x^* C$  by  $f^{-1} : T_{f(x)} C \longrightarrow T_x C$ .

For the rest of this section  $C$  will be hyperelliptic,  $B$  its branched set as in section 3.1 and  $J$  the hyperelliptic involution. This is a central element in  $\text{Aut}(C)$  and we have a short exact sequence

$$0 \longrightarrow \mathbb{Z}_2\{J\} \longrightarrow \text{Aut}(C) \longrightarrow \overline{\text{Aut}}(C) \longrightarrow 0$$

where  $\overline{\text{Aut}}(C)$  is a finite subgroup of  $PSL_2(\mathbb{C})$  and hence is a cyclic, dihedral or polyhedral group. If  $f : C \rightarrow C$  is an automorphism then we associate  $\bar{f} \in \overline{\text{Aut}}(C)$ . Here  $f$  acts on the ramification (i.e. Weierstrass) points and  $\bar{f}$  on the branch set  $B$ . Since an element of  $PSL_2(\mathbb{C})$  other than the identity fixes at most two points, we see that if  $f \neq id, J$ , then  $\bar{f}$  acts on  $B = \{e_1, \dots, e_{2g+2}\}$  with at most two fixed points.

We will be then distinguishing three cases: when  $\bar{f}$  acts fixed point freely, with one fixed point or with two.

We assume below that  $f \neq id, J$ ,  $f$  has order  $n$  so that (wlog)  $\bar{f}(e) = \zeta e$ ,  $\zeta = e^{2\pi i/n}$ . We use the notation  $\langle e \rangle := \{e, \zeta e, \dots, \zeta^{n-1} e\}$  to denote the orbit of  $e \in \mathbb{P}^1(\mathbb{C}) = \mathbb{C} \cup \{\infty\}$  under the action of the cyclic group  $\mathbb{Z}_n \subset PSL_2(\mathbb{C})$  generated by  $\bar{f}$ . Every orbit  $\langle e \rangle$  has length  $n$  except possibly for two which are singletons:  $\langle 0 \rangle = \{0\}$  and  $\langle \infty \rangle = \{\infty\}$ .

A spin structure  $[T]$  as in lemma 3.2 is invariant under the automorphism  $f : C \rightarrow C$  if, and only if,  $\zeta T = T$  or  $T^c$ . We now put this to good use.

**Proposition 3.3.** *Assume  $f \neq id$  has odd order  $n$ . Then the number of invariant spin structures under  $f$  is*

$$\begin{cases} 2^{r-2} & \text{if } \bar{f} \text{ acts freely, where } 2g+2 = nr \\ 2^{r-1} & \text{if } \bar{f} \text{ acts with one fixed point, where } 2g+2 = nr+1 \\ 2^r & \text{if } \bar{f} \text{ acts with two fixed points, where } 2g = nr \end{cases}$$

*Proof.* Since  $n$  is odd an invariant spin structure is determined by a subset  $T \subset B$  such that  $\zeta T = T$  and  $\text{card}(T) \equiv g+1 \pmod{2}$ . Assume  $\bar{f}$  acts freely on  $B$  so that  $B$  is the disjoint union

$$B = \langle e_1 \rangle \sqcup \langle e_2 \rangle \sqcup \dots \sqcup \langle e_r \rangle,$$

where all the  $e_i$  are in  $\mathbb{C}^* = \mathbb{C} - \{0\}$ . By this notation we also mean that we have reordered the  $e_i$  so that  $e_1, \dots, e_r$  are representatives of the orbits  $\langle e_i \rangle$  for  $i = 1, \dots, 2g+2$ . There is no loss of generality in doing this. In that case and if some  $\zeta^m e_i \in T$ , then the entire orbit  $\langle e_i \rangle \subset T$ . Therefore  $T$  must be a disjoint union of the form

$$T = \langle e_{i_1} \rangle \sqcup \langle e_{i_2} \rangle \sqcup \dots \sqcup \langle e_{i_k} \rangle, \text{ where } 1 \leq i_1 < i_2 < \dots < i_k \leq r \text{ and } k = 0, \dots, r.$$

Therefore  $\text{card}(T) = kn \equiv k \pmod{2}$ . The number of such  $T$  is  $\sum_{k=0}^r \binom{r}{k} = 2^r$ , and the number

satisfying  $\text{card}(T) \equiv g+1 \pmod{2}$  is  $2^{r-1}$ . This follows from the identity  $\sum_{k=0}^r (-1)^k \binom{r}{k} = 0$ .

These subsets come in complementary pairs  $\{T, T^c\}$ , and therefore the number of invariant spin structures is  $2^{r-2}$ .

Assume  $\bar{f}$  acts with one fixed point so that

$$B = \langle \cdot \rangle \sqcup \langle e_1 \rangle \sqcup \langle e_2 \rangle \sqcup \dots \sqcup \langle e_r \rangle, \quad e_i \in \mathbb{C}^*, \forall i$$



where  $\langle \cdot \rangle$  is either  $\langle 0 \rangle$  or  $\langle \infty \rangle$ . If  $T \subset B$  satisfies  $\zeta T = T$ , then the possibilities for  $T$  are

$$\begin{aligned} T &= \langle e_{i_1} \rangle \sqcup \langle e_{i_2} \rangle \sqcup \cdots \sqcup \langle e_{i_k} \rangle, \text{ or} \\ T &= \langle \cdot \rangle \sqcup \langle e_{i_1} \rangle \sqcup \langle e_{i_2} \rangle \sqcup \cdots \sqcup \langle e_{i_k} \rangle \end{aligned}$$

In both cases we must have  $1 \leq i_1 < i_2 < \cdots < i_k \leq r$  and  $k = 0, \dots, r$ . The number of such subsets  $T$  is  $2^{r+1}$ , and the number satisfying  $\text{card}(T) \equiv g+1 \pmod{2}$  is  $2^r$ . These subsets come in complementary pairs, and therefore the number of invariant spin structures is  $2^{r-1}$ .

Finally assume that  $\bar{f}$  has two fixed points in  $B$  so that

$$B = \langle 0 \rangle \sqcup \langle \infty \rangle \sqcup \langle e_1 \rangle \sqcup \langle e_2 \rangle \sqcup \cdots \sqcup \langle e_r \rangle, \quad e_i \in \mathbb{C}^*, \forall i$$

If  $T \subset B$  satisfies  $\zeta T = T$  then  $T$  can be any collection of orbits with the right cardinality so that by arguments similar to those above we also get  $2^r$  invariant spin structures.  $\square$

**Proposition 3.4.** *Assume  $f \neq \text{id}$ ,  $J$  has even order  $n$ . Then the number of invariant spin structures is*

$$\begin{cases} 2^{r-1} & \text{if } \bar{f} \text{ acts freely on } B \text{ and } g \text{ is even } (2g+2 = nr) \\ 2^r & \text{if } \bar{f} \text{ acts freely on } B \text{ and } g \text{ is odd } (2g+2 = nr) \\ 2^r & \text{if } \bar{f} \text{ acts on } B \text{ with two fixed points } (2g = nr) \end{cases}$$

Note that when  $n$  is even,  $\bar{f}$  cannot act on  $B$  with a single fixed point since  $2g+2 = nr+1$ .

*Proof.* First assume  $\bar{f}$  acts on  $B$  fixed point freely so that again  $B$  is the disjoint union  $\langle e_1 \rangle \sqcup \langle e_2 \rangle \sqcup \cdots \sqcup \langle e_r \rangle$ , where all the  $e_i$  are in  $\mathbb{C}^* = \mathbb{C} - \{0\}$ . Then there are 2 possibilities for an invariant spin structure  $[T]$ : either  $\zeta T = T^c$  or  $\zeta T = T$ .

If  $\zeta T = T^c$  occurs then necessarily  $T$  is the disjoint union of the sets

$$\{\zeta^{\epsilon_j} e_j, \zeta^{2+\epsilon_j} e_j, \zeta^{4+\epsilon_j} e_j, \dots, \zeta^{n-1+\epsilon_j} e_j\}, \text{ where } j = 1, \dots, r \text{ and } \epsilon_j = \pm 1 \forall j$$

This means that  $n$  is necessarily even (as is the case) and that  $\zeta$  fixes no branch points. The number of such subsets is  $2^r$  since it equals the number of choices of the  $\epsilon_j$ . Every such subset  $T$  determines a spin structure  $[T]$  because  $\text{card}(T) = g+1$  in this case. These subsets come in equivalent pairs  $\{T, T^c\}$ , and therefore there are  $2^{r-1}$  invariant spin structures in this case.

For the second possibility there are  $2^r$  subsets  $T \in \mathcal{S}_B$  such that  $\zeta T = T$ , namely

$$T = \langle e_{i_1} \rangle \sqcup \langle e_{i_2} \rangle \sqcup \cdots \sqcup \langle e_{i_k} \rangle, \text{ where } 1 \leq i_1 < \cdots < i_k \leq r, k = 0, 1, \dots, r$$

Now  $\text{card}(T) = kn \equiv 0 \pmod{2}$ , and therefore  $[T]$  is a spin structure if, and only if,  $g$  is odd. These spin structures come in complementary pairs, so we have  $2^{r-1}$  invariant spin structures for this possibility.

In summary, when  $\bar{f}$  acts fixed point freely there are  $2^{r-1}$  invariant spin structures if  $g$  is even, and  $2^{r-1} + 2^{r-1} = 2^r$  if  $g$  is odd.

The arguments for the other case are similar.  $\square$

**3.3. Genus Two Curves.** As an exercise we count the invariant spin structures for all possible subgroups of  $\text{Aut}(C)$  in the case when  $C$  is of genus two; i.e.  $C : y^2 = f(x)$  where  $f$  is a polynomial of degree 5 or 6. A complete list of all possible reduced automorphism groups that can occur together with their associated equations has been long known (Bolza). For those groups we can count precisely the number of fixed spin structures. It is given by the table below.


Here the actual new statement is in the last column. It is easy to see that the groups in the table are the only ones allowed since besides being finite in  $SO(3)$ , they also are subgroups of  $S_6$  (permutations of the six branch points). Note that the spin structures in genus 2 are given by all equivalence classes of subsets of cardinality 3 in  $B$  (10 of them and these are the even structures) together with all subsets of cardinality 5. For each case in the table above, the configuration of

| $g = 2$ | $\overline{Aut}$ | representative $f$                      | # of fixed spin structures |
|---------|------------------|-----------------------------------------|----------------------------|
| (i)     | $\mathbb{Z}_2$   | $(x^2 - 1)(x^2 - a)(x^2 - b)$           | 4                          |
| (ii)    | $D_4$            | $(x^2 - 1)(x^2 - a)(x^2 - \frac{1}{a})$ | 2                          |
| (iii)   | $D_6$            | $x^6 - 2ax^3 + 1$                       | 1                          |
| (iv)    | $D_{12}$         | $x^6 + 1$                               | 1                          |
| (v)     | $S_4$            | $x(x^4 - 1)$                            | 0                          |
| (vi)    | $\mathbb{Z}_5$   | $x^5 - 1$                               | 1                          |

points of  $B$  in  $\mathbb{P}^1$  is laid out according to the geometry of the associated group action. It is easier to write  $B = \{1, 2, 3, 4, 5, 6\}$ .

*Proof.* (last column of table)

- (i)  $\mathbb{Z}_2$  acts fixed point freely via the map  $x \mapsto -x$  and hence according to proposition 3.4 it fixes  $2^{r-1} = 4$  branch points (here  $n = 2, r = 3$ ).
- (ii) This has to do with the geometry of a parallelogram with vertices at  $a, \bar{a}, -\bar{a}, -a$  say ( $|a| = 1$ ). Let's refer to these branch points by 1, 2, 3, 4 respectively (so that "5" and "6" correspond to the remaining branch points 1 and  $-1$ ). The two involutions are  $\tau_1 = (12)(34)$  (reflection through the  $x$ -axis) and  $\tau_2 = (13)(24)(56)$  (reflection through the  $y$ -axis). According to proposition 3.4,  $\tau_1$  and  $\tau_2$  fix 4 spin structures each but it is easy to see that they have only two such structures in common namely  $\{5, 3, 4\}$  and  $\{5, 1, 2\}$ .
- (iii) This is the situation where the branch points are configured in the plane as in the figure below (each dot represents a branch point in  $\mathbb{C} \subset \mathbb{P}^1$ ).



The polynomial in (iii) has roots  $e^{\pm\alpha}, je^{\pm\alpha}, j^2e^{\pm\alpha}$  numbered as in the figure. There is only one spin structure invariant by the  $\frac{2\pi}{3}$  rotation: namely  $T = \{1, 3, 5\} \sim T' = \{2, 4, 6\}$ . The involution with respect to the  $x$ -axis switches  $T$  and  $T'$  and hence  $D_6$  has a unique invariant spin structure.

- (iv)  $\mathbb{Z}_6$  acts cyclically on  $B$  without fixed points and hence according to proposition 3.4 it has a single fixed spin structure  $T = \{1, 3, 5\} \sim T^c$ .
- (v) the branch points form an octahedron  $\{1, 2, 3, 4, N, S\}$  with north  $N$  and south  $S$  poles. The cyclic  $\mathbb{Z}_4$  acts by rotating the equator  $1 \mapsto 2 \mapsto 3 \mapsto 4$ . According to proposition 3.4 this group fixes 2 spin structures which must be odd of the form  $\{1, 2, 3, 4, N\}$  or  $\{1, 2, 3, 4, S\}$ . But there is an involution switching  $N$  and  $S$  and thus there can't be any invariant spin structure.
- (vi) corresponds to when  $\overline{Aut}$  is of order  $n = 5$  generated by  $\bar{f}$  (in the notation of section 3.2) which acts by fixing one point  $\infty$ . According to proposition 3.3  $\bar{f}$  fixes  $2^{r-1} = 1$  structures.  $\square$

## 4. SPIN STRUCTURES ON KLEIN'S CURVE

In this section we show that the Klein's quartic curve  $\mathcal{K}$  has a unique spin structure invariant under all automorphisms. Let  $G = PSL(2, \mathbf{F}_7) = Aut(\mathcal{K})$ . This is a simple group with presentation of the form

$$G = \langle R, S, T \mid R^2 = S^3 = T^7 = RST = 1, \text{ etc} \rangle$$

Let  $e_1, e_2, \dots, e_6$  be a standard symplectic basis of  $H^1(C; \mathbb{Z})$ . That is a basis of the free abelian group  $H^1(C; \mathbb{Z})$  chosen so that the intersection pairing is given as follows:

$$\langle e_i, e_j \rangle = \begin{cases} +1 & \text{if } j = i + 3 \\ -1 & \text{if } j = i - 3 \\ 0 & \text{in all other cases} \end{cases}$$

From the work of Rauch and Lewittes [14] we can find matrix representations for the induced maps  $R_*, S_*, T_*$  on  $H_1(C; \mathbb{Z})$ :

$$R_* = \begin{bmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & -1 \end{bmatrix}$$

$$S_* = \begin{bmatrix} -1 & 0 & 0 & -1 & -1 & -1 \\ -1 & 0 & 1 & 0 & 0 & -1 \\ -1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 1 \end{bmatrix}$$

$$T_* = \begin{bmatrix} 0 & 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & -1 & -1 & -1 & 0 \\ 0 & 0 & -1 & -1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 1 \\ -1 & 1 & 0 & 0 & -1 & -1 \\ 1 & -1 & 0 & 0 & 1 & 0 \end{bmatrix}$$

The corresponding  $V$  vectors are

$$V_R = [0, 0, 0, 0, 0, 0], V_S = [-1, 1, 1, 0, 0, 0], V_T = [0, 0, 0, -1, 0, 0].$$

Using equation (1), together with a Maple program, it is now direct to determine the sets of spin structures invariant under  $R, S, T$  respectively. The solution vectors  $\bar{X}$  are:

- (1)  $\bar{X} = [x_1, 0, x_3, x_4, x_5, x_4]$ , where the  $x_j$  are arbitrary, for  $R$ -invariant spin structures.
- (2)  $\bar{X} = [x_1 + x_2, x_1 + x_2, 1, x_1 + x_2, x_1, x_2]$ , where the  $x_j$  are arbitrary, for  $S$ -invariant spin structures.
- (3)  $\bar{X} = [0, 0, 1, 0, 0, 0]$  for  $T$ -invariant spin structures.

It is now clear that Klein's curve admits only one invariant spin structure. This is the spin structure fixed by  $T$  and which by choosing the  $x_j$  appropriately we can show is also an invariant for  $R$  and  $S$ .

**4.1. Spin Structures on Quartics.** We would like to say more about what this particular spin structure on  $\mathcal{K}$  is. Generally let  $C$  be a quartic curve in  $\mathbb{P}^2$ . There are  $2^6 = 64$  spin structures on  $C$  as pointed out, of which 28 are “odd” and 36 “even” (see appendix). The canonical divisor class on  $C$  is the locus of the intersection of a hyperplane  $\mathbb{P}^1 \subset \mathbb{P}^2$  with  $C$ . Suppose  $C$  has a bitangent going through the points  $p, q$ ,  $p \neq q$ . This is a line  $\mathbb{P}^1$  intersecting the curve in  $p, q$  with multiplicity two, and hence  $K = 2p + 2q$  (in the divisor class group). By definition,  $p + q$  is a spin structure on  $C$ . Since there are 28 bitangents, this accounts for all odd spin structures.

The even spin structures on the other hand are described as follows. Following [11], a secant line  $\overline{ab}$ ,  $a \neq b \in C$  is a *strict tangent* of  $C$  if either:

- (i)  $\overline{ab}$  is tangent to  $C$  at a point  $r$  different from  $a, b$ , or
- (ii)  $\overline{ab}$  is a triple tangent at  $a$  or at  $b$ .

This is equivalent to the condition that the divisor class  $K = a + b + 2r$  for a point  $r \in C$ . A triangle is now *strictly biscribed* if all sides are strict tangents (see figure 2). A plane cubic has no such triangles but a plane quartic has 288 of them! The way they arise as it turns out is in groups of 8 triangles, one for every even spin structure.




FIGURE 2. A bitangent (odd theta) and a strictly biscribed triangle (even theta)

Let  $\Delta(a, b, c)$  be a strictly biscribed triangle in a quartic  $C$  with double tangents at  $p, q, r$  as in figure 2. Then we can associate to  $\Delta$  the even spin class

$$\theta(\Delta) = a + b + c + p + q + r - K$$

Using the relations  $K = 2p + b + c = 2q + a + c = 2r + a + b$  we can indeed check that  $2\theta(\Delta) = K$ . It is known that for an even theta characteristic  $\theta$ , there are 8 strictly biscribed triangles  $\Delta$  such that  $\theta(\Delta) = \Delta$  [5, 11].

Restricting to the case of the Klein curve  $\mathcal{K}$ , we know that  $\text{Aut}(\mathcal{K}) = \text{PSL}(2, \mathbf{F}_7)$  fixes no bitangent and hence an invariant spin candidate must be even. Consider the points  $a = [1 : 0 : 0]$ ,  $b = [0 : 1 : 0]$  and  $c = [0 : 0 : 1] \in \mathcal{K}$ ,  $\mathcal{K}$  being again the curve  $x^3y + y^3z + z^3x = 0$  in  $\mathbb{P}^2$ . As discussed in [4] for instance, the transformation  $T$  of order 7 can be represented by

$$T : x \mapsto \gamma x, \quad y \mapsto \gamma^4 y, \quad z \mapsto \gamma^2 z$$

where  $\gamma = e^{\frac{2\pi i}{7}}$ . This transformation fixes the three points above and hence the triangle  $xyz = 0$  they define. Note that  $a, b, c$  are inflexion points (i.e. points with triple tangents) and hence they form what Klein calls an inflexion triangle; that is the tangent at  $a$  meets  $\mathcal{K}$  at  $b$ , and the tangent at  $b$  meets  $\mathcal{K}$  at  $c$  and finally the tangent at  $c$  cuts  $\mathcal{K}$  in  $a$ . This is also a biscribed triangle according to our definition with associated theta divisor

$$\theta := a + b + c + a + b + c - K = 2a + 2b + 2c - K$$

(since  $r = a, p = b, q = c$ ).

Let's check that  $\theta$  is fixed by the generators  $S$  and  $R$  of order two and three respectively. Now

$$R : x \mapsto y, \quad y \mapsto z, \quad z \mapsto x$$

so  $\Delta$  and hence  $\theta$  is invariant. To see that  $S$  acts trivially on  $\theta$  it doesn't quite help to write the long expression for  $S$  given also in [4]. Instead consider the quotient  $H = \mathcal{K} / \langle S \rangle$  by the cyclic subgroup generated by  $S$  and write  $\pi : \mathcal{K} \rightarrow H$  the quotient map. Clearly  $H$  cannot be  $\mathbb{P}^1$  since  $\mathcal{K}$  is not hyperelliptic. The only other possibility by the Riemann-Hurwitz formula is that  $\text{genus}(H) = 1$  and  $H$  is an elliptic curve. If  $a \in \mathcal{K}$ , then  $\pi^{-1}(\pi(a)) = \{a, Sa\}$ . But on  $H$  there is a meromorphic function  $f$  with a double pole at  $\pi(a)$  and hence in the divisor class group and by definition  $0 = (f \circ \pi) = 2a - 2S(a)$ . This means that  $2a + 2b + 2c = 2S(a) + 2S(b) + 2S(c)$  and hence  $S(\theta) = \theta$  as desired.

The above discussion proves theorem 1.4.

**Remark 4.1.** In [6], the action of  $PSL(2, \mathbf{F}_7)$  on the set of even spin structures is described with orbit decomposition  $36 = 1 + 7 + 7 + 21$ . The one fixed spin generator is expressed explicitly in terms of the orbital divisors as  $D_2 - D_3 - 7D_7$ , where  $\text{Aut}(C)$  acts on  $\mathcal{K}$  with orbits  $D_2, D_3, D_7$  with stabilizers of orders 2, 3, 7.

## 5. APPENDIX : COHOMOLOGY AND DIVISORS

In this appendix we reconcile both definitions of  $\text{Spin}(C)$  given in §2 and in §3. Let  $Cl(C)$  be the divisor class group of  $C$ ; that is the set of all linearly equivalent divisors. The 0 divisor in  $Cl(C)$  is the divisor of any meromorphic function on  $C$ . The canonical divisor  $K$  is the divisor associated to any meromorphic one form  $\omega$  on  $C$ . We write  $K = (\omega)$ . A spin structure is then a divisor  $D \in Cl(C)$  such that  $2D = K$ . We would like to see that this gives us a cohomology class in  $H^1(U(C); \mathbb{Z}_2)$  which restricts to the generator in the fiber.

The form  $\omega$  is dual to a vector field  $\mathcal{V}$  and the ‘singularities’ of  $\omega$  and  $\mathcal{V}$  are the support of  $D$ ; that is the set of zeros and poles of  $\omega$ . Note that a zero of  $\omega$  is a pole of  $\mathcal{V}$  and vice-versa. If  $(\omega) = 2D$  then the index of the vector field around any of its singular points is even.

Let  $c$  be a closed immersed and framed curve in  $C$  (this is precisely the one cycles in  $H_1(U(C))$ ) and let us assume that  $c$  avoids the singular set of  $\mathcal{V}$ . We can then define the relative winding number of  $\mathcal{V}$  along the curve  $c$  with respect to the framing of  $c$ . This is in words the number of times the vector field rotates in relation to this framing. More specifically, set  $v$  to be the framing, and choose a vector field  $w$  such that  $(v, w)$  is a positively oriented basis along  $c$ . We write  $\mathcal{V} = p_1 v + p_2 w$  where  $p = (p_1, p_2) : S^1 \rightarrow \mathbb{R}^2$ . We then define this winding number to be  $\deg p / |p|$  and we define  $\text{Ind}_{\mathcal{V}}(c) \in \mathbb{Z}_2$  to be this number modulo two. This is a topological invariant and does not change under continuous deformations of the vector field.

Observe at this point that if  $c$  is a 1-boundary (i.e. it bounds a framed disk), then  $\text{Ind}_{\mathcal{V}}(c) = 0$  precisely because the singularities of  $\mathcal{V}$  have even index. On the other hand and if  $c$  is a small simple closed curve with tangential framing, then this framing cannot extend to the interior disk bounded by  $c$ , and consequently the relative winding number is odd and  $\text{Ind}_{\mathcal{V}}(c) = 1$ . Note also that a small closed tangentially framed curve represents the generator of  $H_1(\text{fiber})$  in  $H_1(U(C); \mathbb{Z}_2)$ . The function  $\text{Ind}_{\mathcal{V}}$  is then a well-defined cohomology class

$$\text{Ind}_{\mathcal{V}} \in H^1(U(C), \mathbb{Z}_2) = \text{Hom}(H_1(U(C); \mathbb{Z}_2), \mathbb{Z}_2) = \mathbb{Z}_2^{2g+1}$$

which restricts to the generator of the cohomology of the fibers. This is a spin structure.

If  $c$  is an immersed closed curve in  $C$  that avoids the singularities of  $\omega$ , then the associated quadratic form is given according to [8]

$$q_{\omega}(c) = \text{Ind}_{\mathcal{V}}(c) + \# \text{of components of } c + \# \text{of self-intersections of } c$$

A spin structure  $D \in Cl(C)$  is even (or odd) if the dimension of the vector space of sections of  $L(D)$  the line bundle associated to  $D$  is even (or odd).

ACKNOWLEDGEMENT: The first author is grateful to Jean d'Almeida for interesting conversations regarding [5].

## REFERENCES

- [1] A. Adler; S. Ramanan, *Moduli of abelian varieties*, Lecture Notes in Mathematics, 1644. Springer 1996.
- [2] E. Arbarello, M. Cornalba, P. Griffiths, J. Harris, *Geometry of algebraic curves I*, Springer Grundlehren **267**.
- [3] Atiyah, Michael F. *Riemann surfaces and spin structures*, Ann. scient. École. Norm. Sup.(4) 1971, 47-62.
- [4] C. Bavard, *La surface de Klein*, le journal de maths des élèves de ENS-Lyon (<http://www.ens-lyon.fr/JME/>), vol.1 (1993), 13-22.
- [5] I. Dolgachev, *Topics in classical algebraic geometry, part I*, April 20, 2006.
- [6] I. Dolgachev, *Invariant stable bundles over modular curves  $X(p)$* , Recent progress in algebra (Taejon/Seoul, 1997), 65-99, Contemp. Math., **224**, Amer. Math. Soc., Providence, RI, 1999.
- [7] Farkas, Hershel M. and Kra, Irwin *Riemann Surfaces*, Graduate Texts in Math, v. 71, 1980, Springer Verlag
- [8] Johnson, Dennis *Spin structures and quadratic forms on surfaces*, J. London Math. Soc. (2) **22** (1980), no. 2, 365-373.
- [9] R.C. Kirby, *The topology of 4-manifolds*, Springer Lectures Notes in Math. **1374**, 1980.
- [10] G. Masbaum, *On representations of spin mapping class groups arising in spin TQFT*, Geometry and physics (Aarhus, 1995), 197-207, Lecture Notes in Pure and Appl. Math., 184, Dekker, New York, 1997.
- [11] S. Mukai, *Plane quartics and Fano threefolds of genus twelve*, The Fano Conference, 563-572, Univ. Torino, Turin, 2004.
- [12] Mumford, D. *Theta characteristics of an algebraic curve*, Ann. Sci. cole Norm. Sup. (4) **4** (1971), 181-192.
- [13] Nielsen, J. *Die Structur periodischer Transformation von Flächen*, Danske Vid. Selsk., Mat.-Fys.Medd 15 (1937), 1-77.
- [14] Rauch, Harry E., Lewittes, J. *The Riemann Surface of Klein with 168 Automorphisms*, Problems in analysis (papers dedicated to Solomon Bochner, 1969), 297-308. Princeton Univ. Press, Princeton, N.J., 1970
- [15] P.L. Sipe, *Roots of the canonical bundle of the universal Teichmüller curve and certain subgroups of the mapping class group*, Math. Ann. **260** (1982), no. 1, 67-92.
- [16] Sjerve, Denis *Canonical Forms for Torsion Matrices*, J. of Pure and Algebra **22** (1981) 103-111.

Sadok Kallel  
Laboratoire Painlevé,  
Université de Lille I, France  
[Sadok.Kallel@math.univ-lille1.fr](mailto:Sadok.Kallel@math.univ-lille1.fr)

Denis Sjerve  
Department of Mathematics,  
University of British Columbia,  
Canada  
[sjer@math.ubc.ca](mailto:sjer@math.ubc.ca)